CHILDRENS ALLERGY RELIEF - cetirizine hydrochloride solution Camber Consumer Care

ACTIVE INGREDIENT(S)

(in each 5 mL) Cetirizine Hydrochloride USP 5mg

PURPOSE

Antihistamine

USE(S)

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

WARNINGS

DO NOT USE

if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

ASK A DOCTOR BEFORE USE IF

you have liver or kidney disease. Your doctor should determine if you need a different dose.

ASK A DOCTOR OR PHARMACIST BEFORE USE IF

you are taking tranquilizers or sedatives.

WHEN USING THIS PRODUCT

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

STOP USE AND ASK A DOCTOR IF

an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- use only with enclosed dosing cup
- find right dose on chart below
- mL = milliliter

adults and children 6 years and	5 mL or 10 mL once daily depending upon severity of	
over	symptoms; do not take more than 10 mL in 24 hours.	
adults 65 years and over	5 mL once daily; do not take more than 5 mL in	
_	24 hours.	
children 2 to under 6 years of 2.5 mL once daily. If needed, dose can be increased to a		
age	maximum of 5 mL once daily or 2.5 mL every	
	12 hours. Do not give more than 5 mL in 24	
	hours	
children under 2 years of age	ask a doctor	
consumers with liver or kidney	ask a doctor	
disease		

Other Information

- store between 20° to 25°C (68° to 77°F)
- \bullet do not use if carton is opened or bottle wrap imprinted "SAFETY SEAL®" is broken or missing
- see top panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, flavors, non crystallizing sorbitol solution, propylene glycol, purified water, sodium benzoate and sucralose.

Questions?

call 1-888-588-1418


Distributed by: Camber Consumer Care, Inc.

Piscataway, NJ 08854, USA.

Mfg. Lic. No.: 22/RR/AP/2001/F/R

PRINCIPAL DISPLAY PANEL

Cetirizine Hydrochloride Oral Solution USP, 1 mg/1 mL-container label



Active ingredient (in each 5 mL)

Cetirizine Hydrochloride USP5 mg....Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need.

Purpose a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

uso

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- use only with enclosed dosing cup
- mL= milliliter
- adults and children 6 years and over: 5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours.
- adults 65 years and over: 5 mL once daily; do not take more than 5 mL in 24 hours.
- Children 2 to under 6 years of age: 2.5 mL once daily. If needed, dose can be increased to a maximum of 5 mL once daily or 2.5 mL every

12 hours. Do not give more than 5 mL in 24 hours.

- children under 2 years of age:ask a doctor
- consumers with liver or kidney disease:

ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)
- do not use if bottle wrap imprinted

"SAFETY SEAL®" is broken or missing Inactive ingredients

anhydrous citric acid, flavors, non crystallizing sorbitol solution, propylene glycol, purified water, sodium benzoate and sucralose

Questions? call 1-888-588-1418

Dosing cup should be washed and left to air dry after each use

Cetirizine Hydrochloride Oral Solution USP, 1 mg/1 mL-carton label



CHILDRENS ALLERGY RELIEF

cetirizine hydrochloride solution

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69230-316
--------------	----------------	--------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CETIRIZINE HYDRO CHLO RIDE (UNII: 640047KTOA) (CETIRIZINE - UNII: YO7261ME24)	CETIRIZINE HYDROCHLORIDE	1 mg in 1 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)			
SORBITOL (UNII: 506T60A25R)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			

Product Characteristics			
Color	YELLOW (Colorless to Yellow)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:69230-316-11	1 in 1 CARTON	03/13/2019	
1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210622	03/13/2019	

Labeler - Camber Consumer Care (079539968)

Establishment			
Name	Address	ID/FEI	Business Operations
Hetero Labs Limited Unit III		676162024	ANALYSIS(69230-316), MANUFACTURE(69230-316)

Revised: 3/2019 Camber Consumer Care